

# ON CURVES COVERED BY THE HERMITIAN CURVE

A. COSSIDENTE, G. KORCHMÁROS, AND F. TORRES

ABSTRACT. For each proper divisor  $d$  of  $(q - \sqrt{q} + 1)$ ,  $q$  being a square power of a prime, maximal curves  $\mathbb{F}_q$ -covered by the Hermitian curve of genus  $\frac{1}{2}(\frac{q-\sqrt{q}+1}{d} - 1)$  are constructed.

## 1. INTRODUCTION

A maximal curve over a finite field  $\mathbb{F}_q$ , is a projective geometrically irreducible non-singular algebraic curve defined over  $\mathbb{F}_q$  whose number of  $\mathbb{F}_q$ -rational points attains the Hasse-Weil upper bound

$$q + 1 + 2g\sqrt{q},$$

where  $g$  is the genus of the curve. These curves were studied in [R-Sti], [Sti-X], [FT], [Geer-VI1] (see also the references therein), [FGT], [FT1], [Geer-VI2], [GT], and [CHKT]. Maximal curves have been intensively studied also in connection with Coding Theory in which such curves play an important role [Go].

The Hermitian curve  $\mathcal{H}$ , namely the plane curve defined by

$$(1.1) \quad Y^{\sqrt{q}}Z + YZ^{\sqrt{q}} = X^{\sqrt{q}+1},$$

is a well-known example of a maximal curve over  $\mathbb{F}_q$ . By a result of Lachaud [La, Proposition 6], any non-singular curve  $\mathbb{F}_q$ -covered by  $\mathcal{H}$  is also maximal: indeed almost all of the known maximal curves arise in this way. Lachaud's result has pointed out two fundamental (and still open) problems, namely the classification problem of maximal curves covered by  $\mathcal{H}$  and the existence problem of maximal curves not covered by  $\mathcal{H}$ .

In this paper, we present a new infinite class of maximal curves  $\mathbb{F}_q$ -covered by  $\mathcal{H}$ , which arises from a cyclic automorphism group of  $\mathcal{H}$  of order  $(q - \sqrt{q} + 1)$ , and contains a curve of genus  $\frac{1}{2}(\frac{q-\sqrt{q}+1}{d} - 1)$  for each proper divisor  $d$  of  $(q - \sqrt{q} + 1)$ . The construction together with some remarkable properties are given in Section 5 and Section 6. Since the  $\mathbb{F}_q$ -invariant linear series  $\mathcal{D} := |(\sqrt{q} + 1)P_0|$  plays an important role in some current research on maximal curves over  $\mathbb{F}_q$ , we have computed its dimension and  $(\mathcal{D}, P)$ -orders for some points  $P$ . These curves are non-classical (for the canonical morphism) whenever  $d \leq (q - 2\sqrt{q} + 1)/(2\sqrt{q} - 1)$ . Moreover, for small values of  $d$ , especially for  $d = 3$  (and  $\sqrt{q} \equiv 2 \pmod{3}$ ) our results are also related to recent investigations on

---

1991 Math. Subj. Class.: Primary 11G, Secondary 14G.

This research was carried out within the activity of GNSAGA of the Italian CNR with the support of the Italian Ministry for Research and Technology. Torres acknowledges support of Cnpq-Brazil.

maximal curves having many rational points, or equivalently, large genus with respect to  $q$ . Such maximal curves are somewhat rare. However their classification is a difficult task and still in progress. The first general result in this direction states that the genus of a maximal curve over  $\mathbb{F}_q$  is at most  $\sqrt{q}(\sqrt{q}-1)/2$  with equality holding if and only if the curve is  $\mathbb{F}_q$ -isomorphic to the Hermitian curve, see [Ih], [R-Sti]. In [FT] it is proved that no maximal curve over  $\mathbb{F}_q$  has genus  $g$  with  $g \in [(\sqrt{q}-1)^2/4, \sqrt{q}(\sqrt{q}-1)/2]$ , a result conjectured in [Sti-X]. From [FGT, Theorem 3.1], the second largest genus of maximal curves over  $\mathbb{F}_q$ ,  $q$  odd, is equal to  $(\sqrt{q}-1)^2/4$ , and such curves are  $\mathbb{F}_q$ -isomorphic to the Artin-Schreier curve  $y^{\sqrt{q}} + y = x^{(\sqrt{q}+1)/2}$ . For  $q$  even the second largest genus is  $\sqrt{q}(\sqrt{q}-2)/4$ , and it seems that curves having such a genus are also unique up to  $\mathbb{F}_q$ -isomorphism (cf. [AT]). Instead, the problem of determining the third large genus seems to be much more involved and heavily dependent on certain arithmetical behaviour of  $q$  as the list after Remark 6.5 suggests. From [FT1, Proposition 2.5], the third large genus, for  $q$  odd, is at most  $(\sqrt{q}-1)(\sqrt{q}-2)/4$ . Good candidates come from the known infinite classes of maximal curves  $\mathbb{F}_q$ -covered by  $\mathcal{H}$ , namely  $\mathcal{F}_2$  and  $\mathcal{H}_3$ , where  $\mathcal{F}_t$  is the Fermat curve  $x^{(\sqrt{q}+1)/t} + y^{(\sqrt{q}+1)/t} + 1 = 0$  and  $\mathcal{H}_t$  the Artin-Schreier curve  $y^{\sqrt{q}} + y = x^{(\sqrt{q}+1)/t}$ ,  $t$  being a proper divisor of  $(\sqrt{q}+1)$ . It is worth mentioning that both  $\mathcal{F}_2$  and  $\mathcal{H}_4$  have genus  $(\sqrt{q}-1)(\sqrt{q}-3)/8$ , but they are not  $\mathbb{F}_q$ -isomorphic, [CHKT]. Also the curve  $\mathcal{H}_t$  has been characterized via the type of Weierstrass semigroups at  $\mathbb{F}_q$ -rational points [FGT, Theorem 2.3]. We conjecture that the candidate for the third largest genus in the case of  $\sqrt{q} \equiv 2 \pmod{3}$  is the maximal curve of genus  $(q - \sqrt{q} - 2)/6$  described in Section 6. This emerges from Section 3 where maximal curves with  $\dim(\mathcal{D}) = 3$  are investigated.

As in previous papers concerning maximal curves [FT], [FGT], [FT1], [GT], and [CHKT] we have used Stöhr-Voloch's approach to the Hasse-Weil bound to carry on our research.

**Conventions.** The word curve means a projective geometrically irreducible algebraic curve defined over a finite field  $\mathbb{F}_q$ . For a curve  $\mathcal{X}$ ,  $\tilde{\mathcal{X}}$  denotes its nonsingular model over  $\mathbb{F}_q$ .

## 2. BACKGROUND

In this section we collect some results concerning Weierstrass Point Theory, Frobenius orders and maximal curves.

**2.1. Weierstrass Point Theory and Frobenius orders.** Here we recall relevant material from Stöhr-Voloch's [SV, §1,§2].

Let  $\mathcal{X}$  be a nonsingular curve of genus  $g$  defined over  $\bar{\mathbb{F}}_q$  equipped with the action of the Frobenius morphism  $\text{Fr}_{\mathcal{X}}$  over  $\mathbb{F}_q$ , and let  $\mathcal{D}$  be a  $g_d^r$  on  $\mathcal{X}$ . Suppose that  $\mathcal{D}$  is defined over  $\mathbb{F}_q$ .

Associated to  $\mathcal{D}$  we have two divisors on  $\mathcal{X}$ , namely the *ramification divisor*  $R = R^{\mathcal{D}}$ , and the  $\mathbb{F}_q$ -*Frobenius divisor*  $S = S^{(\mathcal{D}, q)}$ . These divisors provide a lot of geometrical and arithmetical information on  $\mathcal{X}$ . We recall that the set of Weierstrass points  $\mathcal{W}_{\mathcal{X}}$  of  $\mathcal{X}$  is the support of  $R^{\mathcal{K}}$ , where  $\mathcal{K} = \mathcal{K}_{\mathcal{X}}$  is the canonical linear series on  $\mathcal{X}$ .

For  $P \in \mathcal{X}$  let us denote by  $j_i(P) = j_i^{\mathcal{D}}(P)$  the  $i$ -th  $(\mathcal{D}, P)$ -order, by  $\epsilon_i = \epsilon_i^{\mathcal{D}}$  the  $i$ -th  $\mathcal{D}$ -order ( $i = 0, \dots, r$ ), and by  $\nu_i = \nu_i^{(\mathcal{D}, q)}$  the  $i$ -th  $\mathbb{F}_q$ -Frobenius order of  $\mathcal{D}$  ( $i = 0, \dots, r-1$ ). The following are the main properties of  $R$  and  $S$ . Set  $p := \text{char}(\mathbb{F}_q)$ .

1.  $\deg(R) = (2g - 2) \sum_{i=0}^r \epsilon_i + (r + 1)d$ ;
2.  $j_i(P) \geq \epsilon_i$  for each  $i$  and each  $P$ ;
3.  $v_P(R) \geq \sum_{i=0}^r (j_i(P) - \epsilon_i)$  and equality holds iff  $\det\left(\binom{j_i(P)}{\epsilon_j}\right) \not\equiv 0 \pmod{p}$ ;
4.  $(\nu_i)$  is a subsequence of  $(\epsilon_i)$ ;
5.  $\deg(S) = (2g - 2) \sum_{i=0}^{r-1} \nu_i + (q + r)d$ ;
6. For each  $i$  and for each  $P \in \mathcal{X}(\mathbb{F}_q)$ ,  $\nu_i \leq j_{i+1}(P) - j_1(P)$ ;
7. For each  $P \in \mathcal{X}(\mathbb{F}_q)$ ,  $v_P(S) \geq \sum_{i=0}^{r-1} (j_{i+1}(P) - \nu_i)$  and equality holds iff  $\det\left(\binom{j_{i+1}(P)}{\nu_j}\right) \not\equiv 0 \pmod{p}$ .

Therefore if  $P \in \mathcal{X}(\mathbb{F}_q)$ , 6) and 7) imply

8.  $v_P(S) \geq r j_1(P)$ .

Consequently from 5) and 8) we obtain the main result of [SV], namely

9.  $\#\mathcal{X}(\mathbb{F}_q) \leq \deg(S)/r$ .

**2.2. Maximal curves.** We summarize some results from [FGT] and [FT1]. Let  $\mathcal{X}$  be a maximal curve over  $\mathbb{F}_q$ . The key property on  $\mathcal{X}$  is the following linear equivalence [FGT, Cor. 1.2]

$$(2.1) \quad \sqrt{q}P + \text{Fr}_{\mathcal{X}}(P) \sim (\sqrt{q} + 1)P_0, \quad P \in \mathcal{X}, P_0 \in \mathcal{X}(\mathbb{F}_q).$$

Hence, for  $P_0 \in \mathcal{X}(\mathbb{F}_q)$ ,  $\mathcal{X}$  is equipped with the  $\mathbb{F}_q$ -invariant linear series

$$\mathcal{D}_{\mathcal{X}} := |(\sqrt{q} + 1)P_0|,$$

so that  $\dim(\mathcal{D}_{\mathcal{X}})$  is independent of  $P_0 \in \mathcal{X}(\mathbb{F}_q)$ . We have that  $\dim(\mathcal{D}_{\mathcal{X}}) \geq 2$  [Sti-X, Prop. 1] (see also [FGT, Prop. 1.5(iv)]). Furthermore we have the

**Lemma 2.1.** *Let  $\mathcal{X}$  be a maximal curve of genus  $g$  over  $\mathbb{F}_q$ . The following statements are equivalent*

1.  $\mathcal{X}$  is  $\mathbb{F}_q$ -isomorphic to the Hermitian curve (so that  $g = \sqrt{q}(\sqrt{q} - 1)/2$ );
2.  $g > (\sqrt{q} - 1)^2/4$ ;
3.  $\dim(\mathcal{D}_{\mathcal{X}}) = 2$ .

*Proof.* See [R-Sti] and [FT1, Thm. 2.4]. □

Set  $n + 1 := \dim(\mathcal{D}_{\mathcal{X}})$ ,  $j_i(P) := j_i^{\mathcal{D}_{\mathcal{X}}}(P)$ ,  $\epsilon_i := \epsilon_i^{\mathcal{D}_{\mathcal{X}}}$ ,  $\nu_i := \nu_i^{(\mathcal{D}_{\mathcal{X}}, q)}$  and denote by  $m_i(P)$  the  $i$ -th non-gap at  $P \in \mathcal{X}$ . So from (2.1) and [FT1, §2.3], for each  $P \in \mathcal{X}$  the following

holds:

$$(2.2) \quad m_0(P) = 0 < m_1(P) < \dots < m_n(P) = \sqrt{q}.$$

Moreover,  $m_{n+1}(P) = \sqrt{q} + 1$  if  $P \in \mathcal{X}(\mathbb{F}_q)$ , and  $m_{n+1}(P) = \sqrt{q} - 1$  if  $P \notin \mathcal{W}_{\mathcal{X}}$  and  $n+1 \geq 3$  [FGT, Prop. 1.5(iv)(v)]. The main properties of the  $(\mathcal{D}_{\mathcal{X}}, P)$ -orders,  $\mathcal{D}_{\mathcal{X}}$ -orders and  $\mathbb{F}_q$ -Frobenius orders of  $\mathcal{D}_{\mathcal{X}}$  are the following (see [FGT, §1], [FT1, §2.2]).

1. For each  $P \in \mathcal{X}$ ,  $j_1(P) = 1$  so that  $\epsilon_1 = 1$ ;
2.  $\epsilon_{n+1} = \nu_n = \sqrt{q}$ ;
3. If  $P \in \mathcal{X}(\mathbb{F}_q)$ , then the  $(\mathcal{D}_{\mathcal{X}}, P)$ -orders are  $\sqrt{q} + 1 - m_i(P)$  ( $i = 0, 1, \dots, n+1$ ) so that  $v_P(R^{\mathcal{D}_{\mathcal{X}}}) \geq 1$ ;
4. If  $P \notin \mathcal{X}(\mathbb{F}_q)$ , then  $j_{n+1}(P) = \sqrt{q}$  and there exists  $I = I(P) \in [1, n]$  such that  $\sqrt{q} - m_n(P) < \dots < \sqrt{q} - m_{n-I+1}(P) < j_I(P) < \sqrt{q} - m_{n-I}(P) < \dots < \sqrt{q} - m_0(P)$  are the  $(\mathcal{D}_{\mathcal{X}}, P)$ -orders.
5. In particular if  $P \notin \mathcal{W}_{\mathcal{X}}$ , then  $\tilde{m}_i = m_i(P)$  is independent of  $P$ ; hence  $\sqrt{q} - \tilde{m}_i$  is a  $\mathcal{D}$ -order for  $i = 0, 1, \dots, n$ .
6.  $\nu_i = \sqrt{q} - \tilde{m}_{n-i}$  for  $i = 0, \dots, n$ , so that  $\nu_1 = 1$  whenever  $n+1 \geq 3$ .

### 3. ON MAXIMAL CURVES WITH $\dim(\mathcal{D}_{\mathcal{X}}) = 3$

Let  $\mathcal{X}$  be a maximal curve over  $\mathbb{F}_q$  of genus  $g$ . We keep the notations of the previous section. To study  $\mathcal{X}$ , by Lemma 2.1, we can assume that  $g \leq (\sqrt{q}-1)^2/4$  or equivalently that  $\dim(\mathcal{D}_{\mathcal{X}}) \geq 3$ . This section deals with the case  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ . So the  $\mathcal{D}_{\mathcal{X}}$ -orders are  $\epsilon_0 = 0, \epsilon_1 = 1, \epsilon_2$  and  $\epsilon_3 = \sqrt{q}$ , and the  $\mathbb{F}_q$ -Frobenius orders of  $\mathcal{D}$  are  $\nu_0 = 0, \nu_1 = 1$  and  $\nu_2 = \sqrt{q}$  (cf. §2.2(1)(5)(6)).

We first state a sufficient condition to have  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$  and to compute  $\epsilon_2$ .

**Lemma 3.1.** *Let  $X$  be a maximal curve over  $\mathbb{F}_q$  of genus  $g$  such that*

$$(\sqrt{q}-1)(\sqrt{q}-2)/6 < g \leq (\sqrt{q}-1)^2/4.$$

*Then*

1.  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ .
2.  $\epsilon_2 \leq 3$  and  $\epsilon_2 = 2$  provided that  $p = \text{char}(\mathbb{F}_q) \neq 3$ .

*Proof.* (1) Since  $\sqrt{q}, \sqrt{q} + 1 \in H(P)$  for each  $P \in \mathcal{X}(\mathbb{F}_q)$ , (see §2.2),  $\mathcal{D}_{\mathcal{X}}$  is simple and hence we can apply Castelnuovo's genus bound for curves in projective spaces as given in [FGT, p.34]:  $g$  satisfies

$$2g \leq \begin{cases} \frac{(q-n/2)^2}{n} & \text{if } n \text{ is even,} \\ \frac{(q-n/2)^2 - 1/4}{n} & \text{otherwise,} \end{cases}$$

where  $n+1 = \dim(\mathcal{D}_{\mathcal{X}})$ . Therefore  $n+1 \geq 4$  would imply  $g \leq (q-1)(q-2)/6$ , a contradiction. Then we have  $n+1 \leq 3$  and Lemma 2.1 implies  $n+1 = 3$ .

(2) Let  $P \in \mathcal{X}(\mathbb{F}_q)$ . By §2.1(7)(2),  $v_P(S) \geq j_2(P) + 1 \geq \epsilon_2 + 1$ . This inequality, the maximality of  $\mathcal{X}$  and §2.1(5) imply

$$(*) \quad (\sqrt{q} + 1)(2g - 2) + (q + 3)(\sqrt{q} + 1) \geq (\epsilon_2 + 1)[(\sqrt{q} + 1)^2 + \sqrt{q}(2g - 2)].$$

If  $\epsilon_2 \geq 4$ , then we would have

$$(\sqrt{q} + 1)(q - 5\sqrt{q} - 2) \geq (2g - 2)(4\sqrt{q} - 1),$$

and from the upper bound on  $g$  it follows that  $0 > q - 2\sqrt{q} + 10$ , a contradiction.

Now  $\epsilon_2 = 3$  implies  $\binom{3}{2} \equiv 0 \pmod{p}$  by the  $p$ -adic criterion [SV, Cor. 1.9] and hence  $p = 3$ . This finishes the proof of (2).  $\square$

*Remark 3.2.* Here we show that the hypothesis on the genus in Lemma 3.1(1) is sharp whenever  $\sqrt{q} \equiv 0 \pmod{3}$ .

Let  $q = p^m$  with  $m$  even and  $1 \leq r \leq m/2$ . van der Geer and van der Vlugt [Geer-Vl1, Thm. 3.1], [Geer-Vl2, Remark 5.2] constructed a maximal curve  $\mathcal{X}$  over  $\mathbb{F}_q$  of genus  $(p^r - 1)\sqrt{q}/2$ , by considering fibre products of curves of type  $y^p - y = ax^{\sqrt{q}+1}$  with  $a \in \mathbb{F}_q^*$  satisfying  $a^{\sqrt{q}} + a = 0$ . It is not difficult to see that a plane model for  $\mathcal{X}$  is given by an equation of type

$$\sum_{i=0}^r y_1^{p^i} = bx^{\sqrt{q}+1},$$

where  $b \in \mathbb{F}_q^*$  and  $b^{\sqrt{q}} + b = 0$  so that  $\mathcal{X}$  is  $\mathbb{F}_q$ -covered by the Hermitian curve. (The case  $r = m/2$  has worked out in [G-Sti, §V, Example E]).

Set  $\mathcal{D} := \mathcal{D}_{\mathcal{X}}$  and let  $P_0 \in \mathcal{X}(\mathbb{F}_q)$  be the unique point over  $x = \infty$ . Now we can apply the case “ $mn = \ell$ ” in [FT1] to show the following facts:

1.  $\dim(\mathcal{D}) = p^{m/2-r} + 1$ ;
2. The  $(\mathcal{D}, P)$ -orders are as follows:
  - (i)  $0, \sqrt{q} + 1 - ip^r, i = 0, 1, \dots, p^{m/2-r}$  if  $P = P_0$ ;
  - (ii)  $0, 1, \dots, p^{m/2-r}$  and  $\sqrt{q} + 1$  if  $P \in \mathcal{X}(\mathbb{F}_q) \setminus \{P_0\}$ ;
  - (iii)  $0, 1, \dots, p^{m/2-r}$  and  $\sqrt{q}$  if  $P \in \mathcal{X} \setminus \mathcal{X}(\mathbb{F}_q)$ .

In particular, (putting  $p = 3$  and  $r = m/2 - 1$ ) there exists a maximal curve  $\mathcal{X}$  of genus  $\sqrt{q}(\sqrt{q} - 3)/6$  such that  $\dim(\mathcal{D}_{\mathcal{X}}) = 4$ . This shows the sharpness of the upper bound for the genus in Lemma 3.1(1) for  $\sqrt{q} \equiv 0 \pmod{3}$ .

*Remark 3.3.* Let  $\mathcal{X}$  be a maximal curve over  $\mathbb{F}_q$  of genus  $g$  such that  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ .

- (1) If  $\epsilon_2 = 2$ , then §2.1(1), §2.2(3) and the maximality of  $\mathcal{X}$  imply  $g \geq (q - 2\sqrt{q} + 3)/6$ .
- (2) If  $\epsilon_2 = 3$ , then relation  $(*)$  in the proof of Lemma 3.1 implies  $2g - 2 \leq (\sqrt{q} + 1)(q - 4\sqrt{q} - 1)/(3\sqrt{q} - 1)$ . Notice that the upper bound on  $g$  in Lemma 3.1 is equivalent to  $2g - 2 \leq (\sqrt{q} + 1)(\sqrt{q} - 3)/2$ .

Next we investigate the values of  $m_1(P)$  for  $P \in \mathcal{X}(\mathbb{F}_q)$  when  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ . Notice that  $m_1(P) \geq \sqrt{q}/2$  since  $2m_1(P) \in H(P)$  and  $m_2(P) = \sqrt{q}$ , and that  $g$  is bounded from above by the genus of the semigroup  $\langle m_1(P), \sqrt{q}, \sqrt{q} + 1 \rangle$ . We have the following result due to Fuhrmann (see also [Sel, §3.II]).

**Lemma 3.4** (F, §A2). *Let  $\ell, m \in \mathbb{N}$  such that  $m/2 \leq \ell < m$ . Let  $\tilde{g}$  be the genus of the semigroup  $\langle \ell, m, m + 1 \rangle$ .*

1. *If  $\ell \notin \{\lfloor \frac{m+1}{2} \rfloor, m-1, \lfloor \frac{2m+2}{3} \rfloor, m-2\}$ , then*

$$\tilde{g} \leq \begin{cases} \max((m^2 + 4)/8, (m^2 + 3m)/10) & \text{if } \ell \leq 3m/5, \\ (m^2 - 5m + 24)/6 & \text{if } 3m/5 \leq \ell < \lfloor \frac{2m+2}{3} \rfloor, \\ (m^2 - 7m + 70)/6 & \text{if } m \equiv 2 \pmod{3} \text{ and} \\ & (2m + 5)/3 \leq \ell \leq m + 1 - \sqrt{m+1}, \\ (m^2 - 5m + 40)/6 & \text{if } m \equiv 1 \pmod{3} \text{ and} \\ & (2m + 4)/3 \leq \ell \leq m + 1 - \sqrt{m+1}, \\ (m^2 - 3m + 18)/6 & \text{if } \sqrt{q} \equiv 0 \pmod{3} \text{ and} \\ & (2m + 3)/3 \leq \ell \leq m + 1 - \sqrt{m+1}, \\ (m^2 + 2m + 9)/8 & \text{if } m + 1 - \sqrt{m+1} < \ell < m - 2. \end{cases}$$

2. *If  $\ell \in \{\lfloor \frac{m+1}{2} \rfloor, m-1\}$ , then  $\tilde{g} = (m-1)^2/4$ .*

3. *If  $\ell \in \{\lfloor \frac{2m+2}{3} \rfloor, m-2\}$ , then*

$$\tilde{g} = \begin{cases} (m^2 - m + 4)/6 & \text{for } m \equiv 2 \pmod{3}, \\ (m^2 - m)/6 & \text{otherwise.} \end{cases}$$

**Corollary 3.5.** *Let  $\mathcal{X}$  be maximal curve over  $\mathbb{F}_q$ . Suppose that  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$  and that  $\epsilon_2 = 2$ . Then for  $P \in \mathcal{X}(\mathbb{F}_q)$  we have that*

$$m_1(P) \in \{\lfloor \frac{\sqrt{q}+1}{2} \rfloor, \sqrt{q} - 1, \lfloor \frac{2\sqrt{q}+2}{3} \rfloor, \sqrt{q} - 2\}.$$

*Proof.* The genus  $g$  of  $\mathcal{X}$  is bounded from above by the genus of the semigroup  $\langle m_1(P), \sqrt{q}, \sqrt{q} + 1 \rangle$ . Then the result follows by applying the lemma with  $m = \sqrt{q}$  taking into consideration that  $g \geq (q - 2\sqrt{q} + 3)/6$  (Remark 3.3(1)).  $\square$

**Remark 3.6.** Let  $\mathcal{X}$  be a maximal curve over  $\mathbb{F}_q$  and let  $P \in \mathcal{X}(\mathbb{F}_q)$ .

(1) If  $q$  is odd and  $m_1(P) = (\sqrt{q} + 1)/2$ , then  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ ,  $\epsilon_2 = 2$  and  $\mathcal{X}$  is  $\mathbb{F}_q$ -isomorphic to the nonsingular model of  $y^{\sqrt{q}} + y = x^{\sqrt{q}+1}$  [FGT, Thm. 2.3].

(2) If  $q$  is even and  $m_1(P) = \sqrt{q}/2$ , then  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ ,  $\epsilon_2 = 2$  and  $\mathcal{X}$  is  $\mathbb{F}_q$ -isomorphic to the nonsingular model of  $\sum_{i=0}^t y^{\sqrt{q}/2^i} = x^{\sqrt{q}+1}$ ,  $\sqrt{q} = 2^t$  [AT].

**Lemma 3.7.** *Let  $\mathcal{X}$  be a maximal curve over  $\mathbb{F}_q$  such that  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ . Then there exists  $P \in \mathcal{X}(\mathbb{F}_q)$  such that  $m_1(P) + \epsilon_2 = \sqrt{q} + 1$ .*

*Proof.* This follows from the proof of [FGT, Prop. 1.5(v)]; for the sake of completeness we write a proof. By §2.2(3), for each  $P \in \mathcal{X}(\mathbb{F}_q)$ ,  $j_2(P) = \sqrt{q} + 1 - m_1(P)$  so it will be enough to show that  $j_2(P) = \epsilon_2$  for some  $\mathbb{F}_q$ -rational point. Suppose that  $j_2(P) > \epsilon_2$  for each  $P \in \mathcal{X}(\mathbb{F}_q)$  so that  $v_P(R^{\mathcal{D}_X}) \geq 2$  (see §2.1(2)(3)). Using  $\epsilon_2 \leq \sqrt{q} - 1$  and the maximality of  $\mathcal{X}$  from §2.1(1) we have

$$2\sqrt{q}(2g - 2) + 4(\sqrt{q} + 1) \geq 2\sqrt{q}(2g - 2) + 2(\sqrt{q} + 1)^2,$$

i.e.  $0 \geq (\sqrt{q} + 1)(2\sqrt{q} - 2)$ , a contradiction.  $\square$

It follows immediately the

**Corollary 3.8.** *Let  $\mathcal{X}$  be a maximal curve over  $\mathbb{F}_q$  with  $\dim(\mathcal{D}_X) = 3$ . Then the number  $(\sqrt{q} - 2)$  (resp.  $(\sqrt{q} - 1)$ ) is realized as a non-gap at a  $\mathbb{F}_q$ -rational point of  $\mathcal{X}$  iff  $\epsilon_2 = 3$  (resp.  $\epsilon_2 = 2$ ).*

*Remark 3.9.* From Remark 3.6, the number  $\sqrt{q} - 1$  is always realized as a non-gap at  $\mathbb{F}_q$ -rational points of maximal curves. So far we do not know any example of a maximal curve over  $\mathbb{F}_q$  with  $\dim(\mathcal{D}_X) = 3$  and having a  $\mathbb{F}_q$ -rational point  $P$  with  $m_1(P) \in \{\lfloor \frac{2\sqrt{q}+2}{3} \rfloor, \sqrt{q} - 2\}$ .

*Remark 3.10.* If  $q$  is a square and  $\sqrt{q} \equiv 0 \pmod{3}$ , there exists a maximal curve  $\mathcal{X}$  over  $\mathbb{F}_q$  whose genus is  $\sqrt{q}(\sqrt{q} - 1)/6$ . Indeed this is a particular case of the maximal curves constructed in [Geer-Vl2, Prop. 5.1]. From Lemma 3.1 it follows that  $\dim_{\mathcal{D}_X} = 3$ .

We finish this section giving a characterization of the set  $\text{Supp}(S^{\mathcal{X}})$ . Let  $n + 1 = \dim(\mathcal{D}_X)$ ,  $\mathcal{X}$  being a maximal curve over  $\mathbb{F}_q$ . Let  $P \in \mathcal{X} \setminus \mathcal{X}(\mathbb{F}_q)$ . From the proof of [GT, Thm. 2.1] we see that  $P \in \text{Supp}(S^{\mathcal{D}_X}) \Rightarrow m_1(P) < \sqrt{q} - n + 1$ . As an scholium of (loc. cit.), for  $n + 1 = 3$  the converse also holds:

**Lemma 3.11.** *Let  $\mathcal{X}$  be a maximal curve over  $\mathbb{F}_q$  with  $\dim(\mathcal{X}) = 3$ . Then*

$$\text{Supp}(S^{\mathcal{D}_X}) = \mathcal{X}(\mathbb{F}_q) \cup \{P \in \mathcal{X} \setminus \mathcal{X}(\mathbb{F}_q) : m_1(P) < \sqrt{q} - 1\}.$$

*Proof.* We already know that  $\mathcal{X}(\mathbb{F}_q) \cup \{P \in \mathcal{X} \setminus \mathcal{X}(\mathbb{F}_q) : m_1(P) < \sqrt{q} - 1\} \subseteq \text{Supp}(S^{\mathcal{D}_X})$  by §2.1(8) and the remark stated before the lemma. Conversely let  $P \in \mathcal{X} \setminus \mathcal{X}(\mathbb{F}_q)$  with  $m_1(P) < \sqrt{q} - 1$ . Then the  $(\mathcal{D}, P)$ -orders are  $j_0 = 0, j_1 = 1, j_2 = q - m_1(P)$  and  $\sqrt{q}$  (see §2.2(4)). Let  $u, v, w \in \bar{\mathbb{F}}_q(\mathcal{X})$  such that  $j_1 P + D_1 = \text{div}(u) + (\sqrt{q} + 1)P_0$  with  $P \notin \text{Supp}(D_1)$ ;  $\text{div}(v) = D_v - m_1(P)P$  with  $P \notin \text{Supp}(D_v)$  and  $\text{div}(w) = \sqrt{q}P + \text{Fr}_{\mathcal{X}}(P) - (\sqrt{q} + 1)P_0$  (cf. (2.1)). Then

$$\text{div}(uv) + (\sqrt{q} + 1)P_0 = D_u + j_2(P)P + \text{Fr}_{\mathcal{X}}(P) \text{ and } \text{div}(v) + (\sqrt{q} + 1)P_0 = \sqrt{q}P + \text{Fr}_{\mathcal{X}}(P)$$

so, according to the proof of [SV, Thm. 1.1],  $\text{Fr}_{\mathcal{X}}(P)$  belongs to the tangent line at  $P$ . This line is generated by  $\pi(P)$  and  $(D_t^1 \pi)(P)$ , where  $\pi = (1 : u : uv : w)$ ,  $D_t^1 \pi = (0 : D_t^1 u : D_t^1(uv) : D_t^1 w)$  and  $t$  is a local parameter at  $P$ . Since the  $\mathbb{F}_q$ -orders are 0, 1 and  $\sqrt{q}$  (see §2.2(6)) it follows that  $P \in \text{Supp}(S^{\mathcal{D}_X})$ .  $\square$

## 4. PLANE MODELS FOR THE HERMITIAN CURVE

The aim of this section is to introduce two further plane models for the Hermitian curve  $\mathcal{H}$  defined by (1.1).

**4.1. A plane singular model for  $p := \text{char}(\mathbb{F}_q) \geq 3$ .** The Hermitian curve has a cyclic automorphism  $\psi$  of order  $(q - \sqrt{q} + 1)$ . Using (1.1), this automorphism becomes a linear collineation. Unfortunately, the associated matrix is not in diagonal form, and the model (1.1) is not appropriate to investigate those properties of the Hermitian curve which depend on  $\psi$ . This was recognized at first in [C], [CK], [CK1] in the study of  $\psi$ -invariant arcs, or, equivalently, of arcs that are the complete intersection of two Hermitian curves in a suitable mutual position. In [CK], a useful singular plane model was considered for the Hermitian curve. Actually, this model arises from the algebraic envelope of a  $\psi$ -invariant arc, via Segre's fundamental theorem of  $k$ -arcs, see [Se], [H, Thm. 10.4.3]. The same model allows us to determine the quotient curve with respect to the automorphism  $\psi^{(q-\sqrt{q}+1)/d}$ , for each proper divisor  $d$  of  $(q - \sqrt{q} + 1)$ . These curves are maximal and are investigated in §5. Here we limit ourselves to describe how an equation for this model can be obtained. The starting point is the following lemma which follows from the main result in [CK] and [R-Sti].

**Lemma 4.1.** *The nonsingular model  $\bar{\mathcal{C}}$  of the algebraic envelope  $\mathcal{C}$  of a  $\psi$ -invariant arc in  $\mathbb{P}^2(\mathbb{F}_q)$ , regarded as a curve of degree  $2(\sqrt{q} + 1)$  in the dual plane of  $\mathbb{F}_q$ , is  $\mathbb{F}_q$ -isomorphic to the Hermitian curve over  $\mathbb{F}_q$ .*

Let

$$\alpha : \mathbb{P}^2(\bar{\mathbb{F}}_q) \rightarrow \mathbb{P}^2(\bar{\mathbb{F}}_q), \quad (x : y : z) \mapsto (ax : a^{q+1}y : z),$$

where  $a \in \mathbb{F}_{q^3}$  is a primitive  $(q^2 + q + 1)$ -th root of unity, and let

$$E := (1 : 1 : 1).$$

Then the orbit of  $E$  under  $\alpha$  is given by

$$\Pi = \{(a^i : a^{(q+1)i} : 1) : i = 0, 1, \dots, q^2 + q\}.$$

**Lemma 4.2** (CK, Prop. 1). *The set  $\Pi$  is a projective subplane of  $\mathbb{P}^2(\mathbb{F}_{q^3})$  lying in a non-classical position, i.e.  $\Pi \neq \mathbb{P}^2(\mathbb{F}_q)$ .*

Then  $\Pi = \mathbb{P}(\mathbb{L}_q^2)$  with  $\mathbb{L}_q$  a field isomorphic to  $\mathbb{F}_q$ . Now if  $\mathcal{C} : F(X, Y, Z) = 0$ , with  $F \in \mathbb{F}_q[X, Y, Z]$ , one writes the corresponding curve  $\mathcal{C}'$  over  $\bar{\mathbb{L}}_q = \bar{\mathbb{F}}_{q^3}$  as follows. Let

$$A_1 := (1 : 0 : 0), \quad A_2 := (0 : 1 : 0), \quad A_3 := (0 : 0 : 1),$$

and choose three points  $A'_1, A'_2, A'_3 \in \Pi$  such that  $A'_1 A'_2 A'_3 E$  is a non-degenerate quadrangle. Let

$$\kappa : \mathbb{P}^2(\bar{\mathbb{F}}_q) \rightarrow \mathbb{P}^2(\bar{\mathbb{F}}_{q^3})$$



be the projective map such that  $\kappa(A_i) = A'_i$  ( $i = 1, 2, 3$ ) and  $\kappa(E) = E$ . Then  $\mathcal{C}' := \kappa(\mathcal{C})$ . Let

$$\beta : \mathbb{P}^2(\bar{\mathbb{F}}_q) \rightarrow \mathbb{P}^2(\bar{\mathbb{F}}_q), \quad (x : y : z) \mapsto (z : x : y),$$

and

$$\gamma : \mathbb{P}^2(\bar{\mathbb{F}}_q) \rightarrow \mathbb{P}^2(\bar{\mathbb{F}}_q), \quad (x : y : z) \mapsto (x^q : y^q : z^q).$$

**Lemma 4.3.** 1.  $\Pi$  is the set of fixed points of  $\beta \circ \gamma$ , i.e.  $\beta \circ \gamma$  is the Frobenius morphism in the new frame  $A'_1 A'_2 A'_3 E$ .

2. The curve  $\mathcal{C}'$  is defined over  $\mathbb{L}_q$  iff  $\beta(\mathcal{C}) = \mathcal{C}$ .

*Proof.* (1) It is clear that  $\Pi$  is contained in the set of fixed points of  $\beta \circ \gamma$ . Conversely let  $P = (b : c : 1)$  be a fixed point of  $\beta \circ \gamma$ ; then  $(1 : b^q : c^q) = (b : c : 1)$  so that  $bc^q = 1$  and  $b^{q+1} = c$ . Then  $b^{q^2+b+1} = 1$  and (1) follows.

(2) Let  $G = G(X', Y', Z') = 0$  be the equation of  $\mathcal{C}'$ , i.e.  $F = G \circ \kappa$ . Then  $\mathcal{C}'$  is defined over  $\mathbb{L}_q$  iff  $(G \circ \kappa) \circ (\beta \circ \gamma) = \text{Fr} \circ G \circ \kappa$  where  $\text{Fr}$  is the Frobenius morphism on  $\mathbb{P}^1(\bar{\mathbb{L}}_q)$ . So (2) follows from (\*) and the fact that  $\mathcal{C}$  is defined over  $\mathbb{F}_q$ .  $\square$

**Lemma 4.4.** 1. The curve  $\mathcal{C}'$  in coordinates  $(X_0, X_1, X_2) = k(X, Y, Z)$  is defined by

$$(4.1) \quad \begin{aligned} G(X_0, X_1, X_2) = & X_1^2 X_2^{2\sqrt{q}} + X_0^2 X_1^{2\sqrt{q}} + X_0^{2\sqrt{q}} X_2^2 - \\ & 2(X_0^{\sqrt{q}+1} X_1^{\sqrt{q}} X_2 + X_0^{\sqrt{q}} X_1 X_2^{\sqrt{q}+1} + X_0 X_1^{\sqrt{q}+1} X_2^{\sqrt{q}}) = 0. \end{aligned}$$

2. The curve  $\mathcal{C}'$  is defined over  $\mathbb{L}_q$ .

*Proof.* (1) See [CK, Prop. 6].

(2) We know that  $\mathcal{C}$  is defined over  $\mathbb{F}_q$  and that  $F = G \circ \kappa$ . Then to apply Lemma 4.4 it is enough to show that  $G \circ \kappa \circ \beta = G \circ \kappa$ . This easily can be checked using (4.1).  $\square$

Starting from (4.1) we can write an equation for  $\mathcal{C}$  over  $\mathbb{F}_q$  as follows. Let

$$A'_1 = (a : a^{q+1} : 1) \quad A'_2 = (1 : a : a^{q+1}) \quad A'_3 = (a^{q+1} : 1 : a),$$

where  $a \in \mathbb{F}_{q^3}$  is as above. Since  $a^{-q-1} = a^{q^2}$  and  $a^{-1} = a^{q^2+q}$ , then  $A'_2, A'_3 \in \Pi$ . Let  $\kappa$  be the projective map induced by the matrix

$$(4.2) \quad M = \begin{bmatrix} a & 1 & a^{q+1} \\ a^{q+1} & a & 1 \\ 1 & a^{q+1} & a \end{bmatrix}.$$

This map sends  $A_i$  to  $A'_i$  for  $i = 1, 2, 3$  and fixes  $E$ , and since  $(a+1)\det(M) = a^2 + a + 1$ ,  $M$  is non-singular. Hence we have the main result of this subsection:

**Proposition 4.5.** A plane model over  $\mathbb{F}_q$  for the Hermitian curve  $\mathcal{H}$  is given by

$$H(X, Y, Z) = G(aX + Y + a^{q+1}Z, a^{q+1}X + aY + Z, X + a^{q+1}Y + aZ) = 0,$$

with  $G$  defined in (4.1).

**4.2. A plane non-singular model over  $\mathbb{F}_{\sqrt{q^3}}$ .** It is well known that the Hermitian curve  $\mathcal{H}$  is projectively equivalent over  $\mathbb{F}_q$  to the curve of equation

$$H(X, Y, Z) = X^{\sqrt{q}+1} + Y^{\sqrt{q}+1} + Z^{\sqrt{q}+1}.$$

**Proposition 4.6.** *The Hermitian curve  $\mathcal{H}$  is  $\mathbb{F}_{\sqrt{q^3}}$ -isomorphic to the projective plane curve  $\mathcal{C}'$  defined by*

$$G(X_0, X_1, X_2) = X_0^{\sqrt{q}} X_2 + X_2^{\sqrt{q}} X_1 + X_1^{\sqrt{q}} X_0.$$

We first prove the

**Lemma 4.7.** *There exists  $a \in \mathbb{F}_{\sqrt{q^3}}$  satisfying the following properties:*

1.  $a_1 := a^{q\sqrt{q}+\sqrt{q}} + a^{q+\sqrt{q}+1} + a = 0$ ;
2.  $a_2 := a^{q\sqrt{q}+q+\sqrt{q}+1} + a^{\sqrt{q}+1} + 1 = 0$ ;
3.  $a_3 := a^{q\sqrt{q}+\sqrt{q}+1} + a^{q+1} + a^{\sqrt{q}} \neq 0$ ;
4. *The matrix  $M$  in (4.2) is nonsingular.*

*Proof.* We claim that a root  $a$  of the polynomial  $f(X) = X^{\sqrt{q}+1} + X + 1$  satisfies the lemma. In fact, from  $a^{\sqrt{q}+1} + a + 1 = 0$  (\*) we have  $a^{q+\sqrt{q}} + a^{\sqrt{q}} + 1 = 0$  (\*1) so that  $a^{q+\sqrt{q}+1} + a^{\sqrt{q}+1} + a = 0$  (\*2). From (\*) and (\*2),  $a^{q+\sqrt{q}+1} = 1$  (\*3) so that  $a^{\sqrt{q}^3} = a$  (\*4), i.e.  $a \in \mathbb{F}_{\sqrt{q^3}}$ .

Now  $a_1 = a^{q\sqrt{q}+\sqrt{q}} + a^{q+\sqrt{q}+1} + a = a^{1+\sqrt{q}} + a^{q+\sqrt{q}+1} + a = 0$  by (\*4) and (\*1);

$a_2 = a^{q\sqrt{q}+q+\sqrt{q}+1} + a^{\sqrt{q}+1} + 1 = a + a^{\sqrt{q}+1} + 1 = 0$ , by (\*4), (\*3) and (\*).

To prove (3), from (\*4), (\*) and (\*3) we have  $a(a+1)a_3 = -(a^2 + a + 1)^2$  and so  $a_3 = 0$  iff  $a^2 + a + 1 = 0$ . Now  $f(X)$  has  $\sqrt{q} + 1$  different roots; then, as  $\sqrt{q} + 1 \geq 3$ , we can pick a root  $a$  of  $f(X)$  such that  $a^2 + a + 1 \neq 0$ .

Statement (4) follows from the identity  $(a+1)^3 \det(M) = (a^2 + a + 1)^3$  and the proof of (3).  $\square$

*Proof. (Proposition 4.6)* Let  $a \in \mathbb{F}_{\sqrt{q^3}}$  be as in Lemma 4.7. We use the notations of the preceding lemma. Let  $\kappa : \mathbb{P}^2(\bar{\mathbb{F}}_q) \rightarrow \mathbb{P}^2(\bar{\mathbb{F}}_{\sqrt{q^3}})$  be the projective map defined by  $M$  in (4.2). Then  $\kappa^{-1}(\mathcal{C}')$  is the projective plane curve defined by

$$G(aX + Y + a^{q+1}Z, a^{q+1}X + aY + Z, X + a^{q+1}Y + aZ) = 0.$$

After some computations we obtain

$$\begin{aligned} G(aX + Y + a^{q+1}Z, a^{q+1}X + aY + Z, X + a^{q+1}Y + aZ) &= a_3 H(X, Y, Z) + \\ &+ a_2 H(X, Y, Z) + a_1 H(Y, X, Z) \end{aligned}$$

and we are done.  $\square$

5. THE CASE  $d$  DIVIDES  $(q - \sqrt{q} + 1)$ 

Throughout this section we assume that  $q$  is a square power of an odd prime and that  $d$  is a positive divisor of  $(q - \sqrt{q} + 1)$ . The main result here is the

**Theorem 5.1.** *Let  $q$  be a square power of an odd prime. Then for each positive divisor  $d$  of  $(q - \sqrt{q} + 1)$  there exists a maximal curve over  $\mathbb{F}_q$  of genus  $\frac{1}{2}(\frac{q - \sqrt{q} + 1}{d} - 1)$ .*

*Remark 5.2.* If  $\sqrt{q} \equiv r \pmod{d}$  and  $r^2 - r + 1 \equiv 0 \pmod{d}$ , then  $d$  is odd and  $\gcd(r, d) = 1$ . Then  $d = 3$  iff  $r = 2$  and if  $d > 3$ , then 6 divides  $\phi(d)$ ,  $\phi$  being the Euler function. In particular, if  $d$  is prime, then  $d \equiv 1 \pmod{6}$ .

To prove the theorem we use the model  $\mathcal{C}'$  over  $\mathbb{L}_q$  of the Hermitian curve  $\mathcal{H}$  stated in Lemma 4.4. We use the notations of subsection 4.1 and set  $x := X_0/X_2$ ,  $y := X_1/X_2$ . Then, according to (4.1),  $\mathcal{C}'$  is defined by the affine equation

$$y^2 + x^2 y^{2\sqrt{q}} + x^{2\sqrt{q}} - 2(x^{\sqrt{q}+1} y^{\sqrt{q}} + x^{\sqrt{q}} y + x y^{\sqrt{q}+1}) = 0.$$

We recall that  $\mathcal{C}'$  has three singular points, namely  $A_1$ ,  $A_2$  and  $A_3$ , each of them being a 2-fold point and the center of a quadratic branch  $P_i \in \bar{\mathcal{C}}'$ ,  $i = 1, 2, 3$  [CK, Prop. 8]. Moreover if  $t$  is a local parameter at  $P_3$ , a primitive representation of  $P_3$  is given by (loc. cit)

$$(5.1) \quad x = t^2, \quad y = \sum_{i=0}^{\infty} t^{2\sqrt{q}+i(q-\sqrt{q}+1)}.$$

Since  $\beta$  leaves  $\mathcal{C}'$  invariant, maps  $A_3$  into  $A_1$ , and  $A_1$  to  $A_2$  we obtain the following primitive representations for  $P_1$  and  $P_2$  respectively:

$$x = t^{-2\sqrt{q}} + \dots \quad y = t^{-(2\sqrt{q}-2)} + \dots$$

and

$$x = t^{2\sqrt{q}-2} + \dots \quad y = t^{-2} + \dots$$

Hence we have the

**Lemma 5.3.** *The divisors of  $x, y \in \bar{\mathbb{L}}_q(\bar{\mathcal{C}}')$  are respectively:*

$$\operatorname{div}(x) = (2\sqrt{q} - 2)P_2 + 2P_3 - 2\sqrt{q}P_1$$

and

$$\operatorname{div}(y) = 2\sqrt{q}P_3 - (2\sqrt{q} - 2)P_1 - 2P_2.$$

In particular  $x, y \in \mathbb{L}_q(\bar{\mathcal{C}}')$ .

Next consider the morphism

$$\pi = (1 : x^d : y^d) : \bar{\mathcal{C}}' \rightarrow \mathbb{P}^2(\bar{\mathbb{L}}_q).$$

Since  $\pi(P) = (t^{e_P}(P) : (t^{e_P}x^d)(P) : (t^{e_P}y^d)(P))$ , where  $t$  is a local parameter at  $P$  and  $e_P := \min(0, dv_P(x), dv_P(y))$ ,  $v_P$  being the valuation at  $P$ , from the previous lemma we see that  $\pi$  is totally ramified at  $P_1, P_2$  and  $P_3$ . Set  $\mathcal{X}' := \pi(\bar{\mathcal{C}}')$ .

**Lemma 5.4.** *The induced morphism  $\bar{\pi} : \bar{\mathcal{C}}' \rightarrow \bar{\mathcal{X}}'$  is a  $d$ -sheeted covering defined over  $\mathbb{L}_q$  which is ramified precisely at  $P_1, P_2$  and  $P_3$ . Moreover,  $\bar{\pi}$  is totally ramified at each of these points.*

*Proof.* The morphism  $\bar{\pi}$  is defined over  $\mathbb{L}_q$  by Lemma 5.3. From (5.1) we see that  $P'_3 \in \bar{\mathcal{X}}'$  over  $\pi(A_3)$  has a representation of type  $x' = t^{2d}$  and  $y' = (\sum_{i=0}^{\infty} t^{2\sqrt{q}+i(q-\sqrt{q}+1)})^d$ . Putting  $\tau = t^d$  we see that  $P'_3$  has a primitive representation of type

$$x' = \tau^2 \quad y' = \tau^{2\sqrt{q}} + d\tau^{2\sqrt{q}+(q-\sqrt{q}+1)/d} + \dots,$$

showing that  $\bar{\pi}$  has degree  $d$ . Let  $P \in \bar{\mathcal{C}}' \setminus \{P_1, P_2, P_3\}$ , hence a point over  $(a : b : 1) \in \mathcal{C}'$  with  $a \neq 0$  and  $b \neq 0$ . Then  $P$  has a primitive representation of type

$$x = a + t \quad y = b + b_1 t + \dots,$$

with  $b_1 \neq 0$  so that a point  $P' \in \bar{\mathcal{X}}'$  over  $\pi(a : b : 1)$  has a primitive representation of type

$$x' = a^d + dt + \dots \quad y' = b^d + db^{d-1}b_1 t + \dots.$$

This shows that  $\#\bar{\pi}^{-1}(\bar{\pi}(P)) = d$  and the proof is complete.  $\square$

*Proof. (Theorem 5.1)* We show that  $\bar{\mathcal{X}}'$  above, satisfies the theorem. This curve is maximal over  $\mathbb{L}_q$  by Lemma 4.1, Lemma 4.4 and [La, Prop. 6]. To compute the genus  $g$  of  $\bar{\mathcal{X}}'$  we apply the Riemann-Hurwitz formula for  $\bar{\pi} : \bar{\mathcal{C}}' \rightarrow \bar{\mathcal{X}}'$  taking into consideration Lemma 5.4. We have

$$\sqrt{q}(\sqrt{q} - 1) - 2 = d(2g - 2) + 3(d - 1),$$

which gives  $g = \frac{1}{2}(\frac{q-\sqrt{q}+1}{d} - 1)$ .  $\square$

*Remark 5.5.* In the above notations, we have a commutative diagram

$$\begin{array}{ccc} \mathcal{H} = \bar{\mathcal{C}} & \xrightarrow{\kappa} & \mathcal{H} = \bar{\mathcal{C}}' \\ \tilde{\pi} \downarrow & & \downarrow \tilde{\pi} \\ \bar{\mathcal{X}} & \xrightarrow{\kappa} & \bar{\mathcal{X}}' \end{array}$$

where  $\mathcal{X} := \kappa^{-1}(\bar{\mathcal{X}}')$  and  $\tilde{\pi}$  is the morphism induced by  $\kappa^{-1} \circ \pi \circ \kappa$ . We have shown indeed that  $\bar{\mathcal{X}}'$  is a maximal curve over  $\mathbb{L}_q \cong \mathbb{F}_q$ . A plane model for  $\mathcal{X}$  is giving by

$$F(X, Y, Z) = F'(aX + Y + a^{q+1}Z, a^{q+1}X + aY + Z, X + a^{q+1}Y + aZ) = 0,$$

where  $F'(X_0, X_1, X_2) = 0$  is a minimal equation for  $\mathcal{X}'$ . As in the proof of Lemma 4.4, we see that  $\mathcal{X}$  is defined over  $\mathbb{F}_q$  iff  $\beta(\mathcal{X}') = \mathcal{X}'$ . For  $d = 3$  and using another model for  $\mathcal{H}$  we are able to write down an explicit equation for  $\mathcal{X}$  over  $\mathbb{F}_q$  (see next section).

To conclude this section we state some remarks on the dimension of  $\mathcal{D}_{\bar{\mathcal{X}}'} = (\sqrt{q} + 1)P_0$  (cf. §2.2) and on Weierstrass semigroups at some points of  $\bar{\mathcal{X}}'$ . We keep the above notations.

For  $i = 1, 2, 3$ , let  $P'_i \in \bar{\mathcal{X}}'$  be the unique point over  $\pi(A_i)$  and let  $\{Q_i\} := \bar{\pi}^{-1}(P'_i)$ . Since  $\pi(A_1) = A_2$ ,  $\pi(A_2) = A_3$  and  $\pi(A_3) = A_1$  and these points do not belong to  $\Pi$ , then  $Q_1, Q_2$  and  $Q_3$  are not  $\mathbb{L}_q$ -rational. In addition, as  $p$  does not divide  $d$ , the Weierstrass semigroups  $H(Q_i)$  and  $H(P'_i)$  at  $Q_i$  and  $P'_i$  respectively are related to each other as follows (see e.g. [T, proof of Lemma 3.4])

$$S := H(P'_i) = \{h/d : h \in H(Q_i), h \equiv 0 \pmod{d}\}.$$

Moreover,  $H(Q_i)$  can be computed as follows [G-Vi, Thm. 2],

$$\begin{aligned} \tilde{S} := H(Q_i) &= \mathbb{N} \setminus \{r\sqrt{q} + s + 1 : r + s \leq \sqrt{q} - 2\} \\ &= \cup_{j=1}^{\sqrt{q}-2} [j\sqrt{q} - (j-1), j\sqrt{q}] \cup \{0, q - 2\sqrt{q} + 2, q - 2\sqrt{q} + 3, \dots\}. \end{aligned}$$

Hence from (2.2) we have the

**Proposition 5.6.** *For the curve  $\bar{\mathcal{X}}'$  above,*

$$\dim(\mathcal{D}_{\bar{\mathcal{X}}'}) = 1 + \#\{h > 0 : h \in \tilde{S}, h \equiv 0 \pmod{d}, h \leq d\sqrt{q}\}.$$

**Example 5.7.** Let us consider the case  $d = 7$ , i.e.  $7 \mid (q - \sqrt{q} + 1)$  (the case  $d = 3$  is discussed in the next section). By Remark 5.2,  $\sqrt{q} \equiv 3 \pmod{7}$  or  $\sqrt{q} \equiv 5 \pmod{7}$ . The positive elements of  $\tilde{S}$  less than or equal to  $7\sqrt{q}$  are

$$\begin{aligned} &7\sqrt{q} - 6, 7\sqrt{q} - 5, 7\sqrt{q} - 4, 7\sqrt{q} - 3, 7\sqrt{q} - 2, 7\sqrt{q} - 1, 7\sqrt{q} \\ &6\sqrt{q} - 5, 6\sqrt{q} - 4, 6\sqrt{q} - 3, 6\sqrt{q} - 2, 6\sqrt{q} - 1, 6\sqrt{q} \\ &5\sqrt{q} - 4, 5\sqrt{q} - 3, 5\sqrt{q} - 2, 5\sqrt{q} - 1, 5\sqrt{q} \\ &4\sqrt{q} - 3, 4\sqrt{q} - 2, 4\sqrt{q} - 1, 4\sqrt{q} \\ &3\sqrt{q} - 2, 3\sqrt{q} - 1, 3\sqrt{q} \\ &2\sqrt{q} - 1, 2\sqrt{q} \\ &\sqrt{q}. \end{aligned}$$

So we have:

(1) If  $\sqrt{q} \equiv 3 \pmod{7}$ , then  $7\sqrt{q}, 6\sqrt{q} - 4, 5\sqrt{q} - 1$  and  $3\sqrt{q} - 2$  are the elements of  $\tilde{S}$  which are  $\leq 7\sqrt{q}$  and  $\equiv 0 \pmod{7}$ . Thus if  $\sqrt{q} = 3$ , then  $\dim_{\bar{\mathcal{X}}} = 4$ , and if  $\sqrt{q} > 3$ , then  $\dim(\bar{\mathcal{X}}) = 5$ .

(2) If  $\sqrt{q} \equiv 5 \pmod{7}$ , then  $7\sqrt{q}, 6\sqrt{q} - 2, 5\sqrt{q} - 4$  and  $3\sqrt{q} - 1$  are the elements of  $\tilde{S}$  which are  $\leq 7\sqrt{q}$  and  $\equiv 0 \pmod{7}$ . Then  $\dim(\bar{\mathcal{X}}) = 5$ .

Notice that the  $(\mathcal{D}_{\bar{\mathcal{X}}}, P'_i)$ -orders can be computed by means of §2.2(4).

6. THE CASE 3 DIVIDES  $(q - \sqrt{q} + 1)$ 

Throughout this section we let 3 be a divisor of  $(q - \sqrt{q} + 1)$ , or equivalently let  $\sqrt{q} \equiv 2 \pmod{3}$ . We keep the notations of section 4. To construct a curve  $\mathcal{X}$  over  $\mathbb{F}_q$  of genus  $(q - \sqrt{q} - 2)/6$ , we use the equation for  $\mathcal{C}' = \bar{\mathcal{C}}'$  over  $\mathbb{F}_{\sqrt{q}^3}$  given in Proposition 4.6. The construction is similar to the one given in §5 but we work in any characteristic and we shall write down an explicit equation for  $\mathcal{X}$ .

Let  $\mathcal{X}'$  be the curve in  $\mathbb{P}^2(\bar{\mathbb{F}}_q)$  defined by

$$(6.1) \quad \begin{aligned} F'(X_0, X_1, X_2) &:= G(X_0, X_1, X_2) - 3(X_0 X_1 X_2)^{\frac{\sqrt{q}+1}{3}} \\ &= X_0^{\sqrt{q}} X_2 + X_2^{\sqrt{q}} X_1 + X_1^{\sqrt{q}} X_0 - 3(X_0 X_1 X_2)^{\frac{\sqrt{q}+1}{3}}. \end{aligned}$$

and consider the projective morphism

$$\pi = (X_0^3 : X_1^3 : X_2^3) : \mathcal{C}' = \bar{\mathcal{C}}' \rightarrow \mathbb{P}^2(\bar{\mathbb{F}}_q).$$

**Lemma 6.1.** 1. *The curve  $\mathcal{X}'$  is geometrically irreducible.*

2.  $\pi(\mathcal{C}') = \mathcal{X}'$ .

3. *Let  $x := X_0/X_2, y := X_1/X_2 \in \bar{\mathbb{F}}_{\sqrt{q}^3}(\mathcal{C}')$ . Then*

$$\operatorname{div}(x) = (\sqrt{q} - 1)A_2 + A_3 - \sqrt{q}A_1$$

*and*

$$\operatorname{div}(y) = \sqrt{q}A_3 - (\sqrt{q} - 1)A_1 - A_2.$$

4. *The morphism  $\pi$  induces a 3-sheeted covering  $\bar{\pi} : \mathcal{C}' \rightarrow \bar{\mathcal{X}}'$  ramified only at  $A_1, A_2$  and  $A_3$ . Moreover  $\bar{\pi}$  is totally ramified at each of these points.*

5. *The genus of  $\bar{\mathcal{X}}'$  is  $\frac{1}{2}(\frac{q - \sqrt{q} + 1}{3} - 1)$ .*

*Proof.* (1) Each fundamental line  $X_i = 0$  ( $i = 0, 1, 2$ ) meets  $\mathcal{X}'$  in only two points. Assume  $\mathcal{X}'_1$  be a component of  $\mathcal{X}'$ . Clearly  $\mathcal{X}'_1$  must meet each fundamental line in at least one point. This implies that at least two points in the set  $S := \{A_1, A_2, A_3\}$  lie on  $\mathcal{X}'_1$ . If we would had a further component  $\mathcal{X}'_2$  of  $\mathcal{X}'$ , then this component would also pass through two points of  $S$  and one of these points certainly be a common point of  $\mathcal{X}'_1$  and  $\mathcal{X}'_2$ . Then a point of  $S$  would be a singular point of  $\mathcal{X}'$  which is not the case, as a direct computation shows.

(2) From the identity

$$a^3 + b^3 + c^3 - 3abc = (a + b + c)(\epsilon^2 a + b + \epsilon c)(\epsilon a + b + \epsilon^2 c),$$

where  $\epsilon$  is a primitive 3-th root of unity, we have that

$$F'(X_0^3, X_1^3, X_2^3) = G(X_0, X_1, X_2)G(\epsilon X_0, \epsilon X_1, X_2)G(X_0, X_1, \epsilon X_2).$$

This and (1) implies (2).

(3) Similar to the proof of Lemma 5.3.

(4) Similar to the proof of Lemma 5.4.

(5) Follows from (4) and the Riemann-Hurwitz formula applied to  $\bar{\pi}$ .  $\square$

Consider the following commutative diagram

$$\begin{array}{ccc} \mathcal{H} & \xrightarrow{\kappa} & \mathcal{C}' = \bar{\mathcal{C}}' \\ \bar{\pi} \downarrow & & \downarrow \bar{\pi} \\ \bar{\mathcal{X}} & \xrightarrow{\kappa} & \bar{\mathcal{X}}' \end{array}$$

where  $\mathcal{X} := \kappa^{-1}(\mathcal{X}')$  and  $\bar{\pi}$  is the morphism induced by  $\kappa^{-1} \circ \pi \circ \kappa$ . Now we can state the main result of this section.

**Theorem 6.2.** *The curve  $\bar{\mathcal{X}}$  above is a maximal curve over  $\mathbb{F}_q$  of genus  $(q - \sqrt{q} - 2)/6$ . A plane model over  $\mathbb{F}_q$  for  $\mathcal{X}$  is given by*

$$(6.2) \quad F(X, Y, Z) = cF'(aX + Y + a^{\sqrt{q}+1}Z, a^{q+1}X + aY + Z, X + a^{q+1}Y + aZ) = 0,$$

where  $a \in \mathbb{F}_{\sqrt{q}^3}$  satisfies Lemma 4.7,  $c \in \mathbb{F}_{q^3}$  such that  $c^{\sqrt{q}-1} = a$ , and  $F'(X_0, X_1, X_2)$  is the polynomial in (6.1).

*Proof.* The statement on the genus follows from Lemma 6.1(5) and the diagram above. To see that  $\bar{\mathcal{X}}$  is maximal, by [La, Proposition 6], it is enough to show that  $\mathcal{X}$  is defined over  $\mathbb{F}_q$ . Let  $G'$  be a minimal equation of  $\mathcal{X}'$  so that

$$G'(X, Y, Z) := F'(aX + Y + a^{q+1}Z, a^{q+1}X + aY + Z, X + a^{q+1}Y + aZ) = 0.$$

Note that  $F'(X_0, X_1, X_2) \in \mathbb{F}_q[X_0, X_1, X_2]$  and that  $F'(X_0, X_1, X_2) = F'(X_2, X_0, X_1)$ , i.e.  $\beta(\mathcal{X}') = \mathcal{X}'$ . Using these facts we obtain

$$\begin{aligned} G'(X, Y, Z)^q &= F'(a^q X^q + Y^q + a^{q^2+q} Z^q, a^{q^2+q} X^q + a^q Y^q + Z^q, X^q + a^{q^2+q} Y^q + a^q Z^q) \\ &= F'(X^q + a^{q^2+q} Y^q + a^q Z^q, a^q X^q + Y^q + a^{q^2+q} Z^q, a^{q^2+q} X^q + a^q Y^q + Z^q). \end{aligned}$$

Now from  $a^{q+\sqrt{q}+1} = 1$  (cf. proof of Lemma 4.7) we get  $a^{q^2+q+1} = 1$  and since  $F'(X_0, X_1, X_2)$  is a homogeneous polynomial of degree  $\sqrt{q} + 1$  we have that

$$G'(X, Y, Z)^q = a^{-(\sqrt{q}+1)} G'(X^q, Y^q, Z^q).$$

Then  $\mathcal{X}$  can also be defined by

$$F(X, Y, Z) := cG'(X, Y, Z) \quad \text{with} \quad c^{\sqrt{q}-1} = a,$$

(so that  $c \in \mathbb{F}_{q^3}$  as  $a^{q^2+q+1} = 1$ ), and since  $F(X, Y, Z)^q = F(X^q, Y^q, Z^q)$  we are done.  $\square$

Next we investigate the linear system  $\mathcal{D}_{\bar{\mathcal{X}}} := |(\sqrt{q} + 1)P_0|$  (cf. §2.2). For  $i = 1, 2, 3$ , let  $P'_i \in \bar{\mathcal{X}}'$  be the unique point over  $\pi(A_i)$  and let  $P_i := \kappa^{-1}(P'_i) \in \bar{\mathcal{X}}$ . We notice that  $P_i \notin \bar{\mathcal{X}}(\mathbb{F}_q)$  because  $\pi(A_i) \in \{A_1, A_2, A_3\}$  and the coordinates of  $\kappa^{-1}(A_i)$  do not belong to  $\mathbb{F}_q$ .

**Lemma 6.3.** *The first two positive Weierstrass non-gaps at  $P_i$  above ( $i = 1, 2, 3$ ) are  $(2\sqrt{q} - 1)/3$  and  $\sqrt{q}$ .*

*Proof.* Is a particular case of the discussion after Remark 5.5.  $\square$

**Proposition 6.4.** *Let  $\bar{\mathcal{X}}$  be the maximal curve over  $\mathbb{F}_q$  of Theorem 6.2. Then*

1.  $\dim(\mathcal{D}_{\bar{\mathcal{X}}}) = 3$ .
2. *The  $(\mathcal{D}_{\bar{\mathcal{X}}}, P)$ -orders are:*
  - (i)  $0, 1, 2$  and  $\sqrt{q} + 1$  if  $P \in \bar{\mathcal{X}}(\mathbb{F}_q)$ ;
  - (ii)  $0, 1, (\sqrt{q} + 1)/3$  and  $\sqrt{q}$  if  $P \in \{P_1, P_2, P_3\}$ ;
  - (iii)  $0, 1, 2$  and  $\sqrt{q}$  if  $P \in \bar{\mathcal{X}} \setminus (\bar{\mathcal{X}}(\mathbb{F}_q) \cup \{P_1, P_2, P_3\})$ .

*Proof.* Set  $\mathcal{D} := \mathcal{D}_{\bar{\mathcal{X}}}$ .

(1) It follows from Lemma 3.1(1) (or from Prop. 5.6 with  $d = 3$ ).

(2) The  $\mathcal{D}$ -orders are  $0, 1, 2$  and  $\sqrt{q}$  by Lemma 3.1(2). From Lemma 6.3 and §2.2(1)(4) for each  $P_i$  ( $i = 1, 2, 3$ ) the  $(\mathcal{D}, P_i)$ -orders are  $0, 1, (\sqrt{q} + 1)/3$  and  $\sqrt{q}$ . Hence from §2.1(3), it follows that  $v_{P_i}(R^{\mathcal{D}}) = (\sqrt{q} - 5)/3$ . Now, using  $g = (q - \sqrt{q} - 2)/6$  and the maximality of  $\bar{\mathcal{X}}$  we obtain

$$\deg(R^{\mathcal{D}}) - 3(\sqrt{q} - 5)/3 = \#\bar{\mathcal{X}}(\mathbb{F}_q).$$

Then (2) follows by taking into consideration that  $j_3(P) = \sqrt{q} + 1$  at each  $\mathbb{F}_q$ -rational point (cf. §2.2(3)).  $\square$

*Remark 6.5.* We can describe  $\mathcal{D}_{\bar{\mathcal{X}}}$  via hyperplane sections on a certain curve. Indeed, consider the morphism

$$\varphi = (1 : x^3 : y^3 : xy) : \mathcal{C}' \rightarrow \mathbb{P}^3(\bar{\mathbb{F}}_{\sqrt{q}^3})$$

with  $x = X_0/X_2, y = X_1/X_2 \in \bar{\mathbb{F}}_{\sqrt{q}^3}(\mathcal{C}')$  satisfying  $x^{\sqrt{q}} + y + xy^{\sqrt{q}} = 0$ . Let  $\mathcal{Y} := \varphi(\mathcal{C}')$  and let  $\text{Pr}$  denotes the projection of  $\mathbb{P}^3(\bar{\mathbb{F}}_q) \setminus \{(0 : 0 : 0 : 1)\}$  from the point  $(0 : 0 : 0 : 1)$  on the plane  $\mathbb{P}^2(\bar{\mathbb{F}}_q) \cong \{(x : y : z : 0) : (x : y : z) \in \mathbb{P}^2(\bar{\mathbb{F}}_q)\}$ . Then arguing as in Lemma 5.4 one can show that

- (i)  $\mathcal{X}' = \text{Pr}(\mathcal{Y})$ ;
- (ii)  $\mathcal{D} = (\kappa^{-1} \circ \text{Pr})^*(\mathcal{D}_{\bar{\mathcal{X}}})$ ,

where  $\mathcal{D}$  cuts  $\mathcal{Y}$  by hyperplanes of  $\mathbb{P}^3(\bar{\mathbb{F}}_{\sqrt{q}^3})$ .

So far we have the following maximal curves  $\mathcal{X}$  over  $\mathbb{F}_q$  such that  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$ :

1. The non-singular model of  $y^{\sqrt{q}} + y = x^{(\sqrt{q}+1)/2}$ ,  $q$  odd, whose genus is  $(\sqrt{q} - 1)^2/4$  (cf. Remark 3.6(1));
2. The non-singular model of  $\sum_{i=0}^t y^{\sqrt{q}/2^i} = x^{\sqrt{q}+1}$ ,  $q = 2^t$ , whose genus is  $\sqrt{q}(\sqrt{q} - 2)/4$  (cf. Remark 3.6(2));
3. A maximal curve of genus  $\sqrt{q}(\sqrt{q} - 1)/6$ ,  $\sqrt{q} \equiv 0 \pmod{3}$ , (cf. Remark 3.10).
4. A maximal curve of genus  $(q - \sqrt{q} - 2)/6$  (cf. Theorem 6.2).

A natural question then is the following:



**Question.** Are all the maximal curves  $\mathcal{X}$  over  $\mathbb{F}_q$  (up to  $\mathbb{F}_q$ -isomorphism) with  $\dim(\mathcal{D}_{\mathcal{X}}) = 3$  listed above ?

*Remark 6.6.* Connection between maximal curves and non-classical curves have been noticed in [FGT, Prop. 1.7]. More precisely, a maximal curve over  $\mathbb{F}_q$  of genus  $g$  is non-classical provided that  $g \geq \sqrt{q} - 1$ . Hence the examples obtained here are non-classical for  $d \leq (q - \sqrt{q} + 1)/(2\sqrt{q} - 1)$ . Most of the known examples of non-classical curves are Artin-Schreier extensions of rational function fields, with genus a multiple of the characteristic [Sch], [G-Vi]. Thus the maximal curves obtained here are in fact new examples of non-classical curves.

## REFERENCES

- [AT] M. Abdón and F. Torres, On maximal curves in characteristic two, in preparation.
- [C] A. Cossidente, A new proof of the existence of  $(q^2 - q + 1)$ -arcs in  $PG(2, q^2)$ , *J. Geom.* **53** (1995), 37–40.
- [CHKT] A. Cossidente, J.W.P. Hirschfeld, G. Korchmáros and F. Torres, On plane maximal curves, Preprint (1998), (alg-geom/9802113).
- [CK] A. Cossidente and G. Korchmáros, The algebraic envelope associated to a complete arc, *Rend. Circ. Mat. Palermo, Suppl.* **51** (Recent Progress in Geometry, E. Ballico, G. Korchmáros Eds.) (1998), 9–24.
- [CK1] A. Cossidente and G. Korchmáros, The Hermitian function field arising from a cyclic arc in a Galois plane, *Geometry, Combinatorial Designs and Related Structures*, London Math. Soc. Lecture Note Series **245**, Cambridge University Press, Cambridge, 63–68, 1997.
- [F] R. Fuhrmann, *Algebraische Funktionenkörper über endlichen Körpern mit maximaler Anzahl rationaler Stellen*, Ph.D. dissertation, Universität GH Essen, Germany, 1995.
- [FGT] R. Fuhrmann, A. Garcia and F. Torres, On maximal curves, *J. Number Theory* **67**(1) (1997), 29–51.
- [FT] R. Fuhrmann and F. Torres, The genus of curves over finite fields with many rational points, *Manuscripta Math.* **89** (1996), 103–106.
- [FT1] R. Fuhrmann and F. Torres, On Weierstrass points and optimal curves, *Rend. Circ. Mat. Palermo Suppl.* **51** (Recent Progress in Geometry, E. Ballico, G. Korchmáros Eds.) (1998), 25–46.
- [G-Sti] A. Garcia and H. Stichtenoth, Algebraic function fields over finite fields with many places, *IEEE Trans. Inform. Theory* **41**(6) (1995), 1548–1563.
- [GT] A. Garcia and F. Torres, On maximal curves having classical Weierstrass gaps, submitted.
- [G-Vi] A. Garcia and P. Viana, Weierstrass points on certain non-classical curves, *Arch. Math.* **46** (1986), 315–322.
- [Geer-Vl1] G. van der Geer and M. van der Vlugt, How to construct curves over finite fields with many points, *Arithmetic Geometry* (Cortona 1994), F. Catanese Ed., Cambridge Univ. Press, Cambridge, 169–189, 1997.
- [Geer-Vl2] G. van der Geer and M. van der Vlugt, Generalized Reed-Muller codes and curves with many points, Report W97-22, Mathematical Institute, University of Leiden, The Netherlands, (alg-geom/9710016).
- [Go] V.D. Goppa, Algebraic-Geometric codes, *Math. USSR-Izv.* **21** (1), 75–91 (1983).
- [H] J.W.P. Hirschfeld, *Projective Geometries Over Finite Fields*, second edition, Oxford University Press, Oxford, 1998.

- [HK] J.W.P. Hirschfeld and G. Korchmáros, Embedding an arc into a conic in a finite plane, *Finite Fields Appl.* **2** (1996), 274–292.
- [HK1] J.W.P. Hirschfeld and G. Korchmáros, The number of points on an algebraic curve over a finite field, *Bull. Belg. Math. Soc. Simon Stevin*, to appear.
- [Ih] Y. Ihara, Some remarks on the number of rational points of algebraic curves over finite fields, *J. Fac. Sci. Tokio* **28** (1981), 721–724.
- [La] G. Lachaud, Sommes d’Eisenstein et nombre de points de certaines courbes algébriques sur les corps finis, *C.R. Acad. Sci. Paris* **305**, Série I (1987), 729–732.
- [R-Sti] H.G. Rück and H. Stichtenoth, A characterization of Hermitian function fields over finite fields, *J. Reine Angew. Math.* **457** (1994), 185–188.
- [Sch] F.K. Schmidt, Zur arithmetischen Theorie der algebraischen Funktionen II. Allgemeine Theorie der Weierstrasspunkte, *Math. Z.* **45**, (1939), 75–96.
- [Se] B. Segre, Introduction to Galois geometries (edited by J.W.P. Hirschfeld), *Atti Accad. Naz. Lincei* **8** (1967), 133–236.
- [Sel] E.S. Selmer, On the linear diophantine problem of Frobenius, *J. Reine Angew. Math.* **293/294** (1977), 1–17.
- [Sti-X] H. Stichtenoth and C.P. Xing, The genus of maximal function fields, *Manuscripta Math.* **86** (1995), 217–224.
- [SV] K.O. Stöhr and J.F. Voloch, Weierstrass points and curves over finite fields, *Proc. London Math. Soc.* **52** (1986), 1–19.
- [T] F. Torres, On certain  $N$ -sheeted coverings of curves and numerical semigroups which cannot be realized as Weierstrass semigroups, *Comm. Algebra* **23**(11) (1995), 4211–4228.

DIPARTIMENTO DI MATEMATICA UNIVERSITÀ DELLA BASILICATA VIA N. SAURO 85, 85100 POTENZA, ITALY

*E-mail address:* `cossidente@unibas.it`

*E-mail address:* `korchmaros@unibas.it`

IMECC-UNICAMP, Cx. P. 6065, CAMPINAS-13083-970-SP, BRAZIL

*E-mail address:* `ftorres@ime.unicamp.br`